

HEPATITIS B REFERRAL FORM

PATIENT INFORMATION					
Patient Name:		DOB:	Sex: <input type="checkbox"/> M <input type="checkbox"/> F	Weight:	<input type="checkbox"/> lbs. <input type="checkbox"/> kg.
SSN:	Phone:	Allergies:			
Address:		City:	State:	Zip:	
Emergency Contact:		Phone:	<input type="checkbox"/> Please attach demographic information		
PRESCRIBER INFORMATION					
Prescriber:		NPI:	DEA:	State Lic:	
Supervising Physician:		Practice Name:			
Address:		City:	State:	Zip:	
Phone:	Fax:	Key Office Contact:		Phone:	
DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT					
Primary Diagnosis: <input type="checkbox"/> Hepatitis B <input type="checkbox"/> HIV-HBV Co-infection <input type="checkbox"/> Other: _____					
Medical Assessment: Please provide the information below of fax copies of labs to above number.					
PCR for HBV DNA (Viral Load) _____, Date: _____ AST/ALT _____ CrCl _____					
Ratio _____ / _____ Date: _____ <input type="checkbox"/> e-antigen + (HBeAg+) / <input type="checkbox"/> e-antigen - (HBeAg-),					
<ul style="list-style-type: none"> ■ Has patient been treated <i>previously</i> for this condition? <input type="checkbox"/> Yes <input type="checkbox"/> No Medication(s): _____ ■ Is patient <i>currently</i> on therapy? <input type="checkbox"/> Yes <input type="checkbox"/> No Medication(s): _____ ■ Will patient stop taking the above medication(s) before starting the new medication? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes: _____ ■ How long should patient wait before starting the new medication? _____ ■ Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____ 					
INSURANCE INFORMATION					
<input type="checkbox"/> Please attach front and back of patient's insurance card (medical and prescription)					
COPAY CARD ENROLLMENT					
<input type="checkbox"/> Please check if enrolling in copay card		Copay ID: _____			
PRESCRIPTION INFORMATION					
<input type="checkbox"/> Baraclude					
<input type="checkbox"/> 0.5 mg tab: Take 1 tablet PO daily on empty stomach (Naive PT)			QTY: <u>30 Tabs</u>	Refills: _____	
<input type="checkbox"/> 1 mg tab: Take 1 tablet PO daily on empty stomach (Lamivudine-Refractory or decompensated liver disease)			QTY: <u>30 Tabs</u>	Refills: _____	
<input type="checkbox"/> 0.05 mg/mL oral solution: _____			QTY: _____	Refills: _____	
<input type="checkbox"/> Alternate Dose (CrCL<50 mL/min or Dialysis): _____			QTY: _____	Refills: _____	
<input type="checkbox"/> Epivir HBV 100 mg					
<input type="checkbox"/> 100 mg PO daily			QTY: <u>30 Tabs</u>	Refills: _____	
<input type="checkbox"/> Epivir HBV 150 mg					
<input type="checkbox"/> 150 mg PO BID (only for co-infected PT with HIV)			QTY: <u>60 Tabs</u>	Refills: _____	
<input type="checkbox"/> Epivir HBV 5mg/ml oral solution: _____			QTY: _____	Refills: _____	
<input type="checkbox"/> Alternate Dose (CrCL<50 mL/min or Dialysis): _____			QTY: _____	Refills: _____	
<input type="checkbox"/> HBIG Hepatitis B Immune Globulin-single use vial, greater than 1560 International Units/5 mL, greater than 312 International Units/mL					
<input type="checkbox"/> 5 mL IM in 2 divided doses <input type="checkbox"/> Once or <input type="checkbox"/> every 28 days.			QTY: <u>5 mL vial</u>	Refills: _____	
<input type="checkbox"/> 2 mL IM in 2 divided doses <input type="checkbox"/> Once or <input type="checkbox"/> every 28 days.			QTY: <u>2 of 1 mL vial</u>	Refills: _____	
<input type="checkbox"/> HBIG MD Infusion (1560 International Units/5ml vials)					
<input type="checkbox"/> 20,000 International Units (64mL) in 250 mL NS, IV over _____ hour(s), every _____ for _____ infusions			QTY: _____ Vials	Refills: _____	
<input type="checkbox"/> Alternate Dose: _____			QTY: _____ Vials	Refills: _____	
<input type="checkbox"/> Hepsera 10 mg					
<input type="checkbox"/> 10 mg PO daily			QTY: <u>30 Tabs</u>	Refills: _____	
<input type="checkbox"/> Alternate Dose (CrCL<50 mL/min or Dialysis): _____			QTY: _____	Refills: _____	
<input type="checkbox"/> Pegasys 180 mcg Prefilled Syringe (OR) <input type="checkbox"/> Pegasys 180 mcg Vial *Will dispense prefilled syringe unless vial is marked					
<input type="checkbox"/> 180 mcg SQ once every week			QTY: <u>28 days</u>	Refills: _____	
<input type="checkbox"/> Alternate Dose (CrCL<50 mL/min or Dialysis): _____			QTY: _____	Refills: _____	
<input type="checkbox"/> Vemlidy 25 mg					
<input type="checkbox"/> 25 mg PO daily with food			QTY: <u>30 Tabs</u>	Refills: _____	
			QTY: <u>30 Tabs</u>	Refills: _____	
<input type="checkbox"/> Viread 300 mg					
<input type="checkbox"/> 300 mg PO daily			QTY: <u>30 Tabs</u>	Refills: _____	
<input type="checkbox"/> Alternate Dose (CrCL<50 mL/min or Dialysis): _____			QTY: _____	Refills: _____	
<input type="checkbox"/> Other: _____					
			QTY: _____	Refills: _____	

Prescriber's Signature: _____ DAW (Dispense as Written) **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send prescription on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.