

A. Patient information

First name _____ Last name _____
 Address _____
 City _____ State _____ ZIP _____
 Email _____
 Phone (home) _____ Phone (mobile) _____
 Preferred contact: Text Email Phone (home) Phone (mobile)
 Preferred language: English Spanish Other _____
 Sex on file with insurance: Male Female Date of birth: ____/____/____

Prescription insurance information


Attach copies of both sides of patient's pharmacy benefit card(s).

Patient does **not** have coverage (complete the Patient Assistance Application on page 5)
 Patient has secondary insurance

Primary insurance name _____ Policy no. _____
 Group no. _____ Insurance company phone _____
 Policy holder name _____
 Prior authorization number (if known) _____ Prior authorization effective dates _____

Patient authorization

I have read and agree to the Authorization to Use and Disclose Personal Information on Page 2 and the full Terms and Conditions on Page 3. You must sign below to participate in Interconnect Support Services.

Signature of patient/legal representative _____ Date _____/_____/_____


Name of patient/legal representative _____

Legal representative phone _____ Relationship to patient _____

Additional opt-in: By checking this box, I authorize Intercept, and third-party companies working on Intercept's behalf, to contact me by mail, email, telephone, or text message for (i) marketing purposes, including to provide me with information about Intercept's products, services, and programs or other topics of interest, and/or (ii) to conduct market research.

I have read and agree to the Terms and Conditions for the Interconnect Savings Program on page 3.

B. Prescriber information

First name _____ Last name _____
 Address _____
 City _____ State _____ ZIP _____
 Phone _____ Ext. _____ Fax _____
 Primary contact name _____ Primary contact phone _____
 Primary contact fax _____ Clinic/facility name _____
 NPI no. _____ State license no. _____ Prescriber tax ID _____
 All correspondence regarding this patient will be communicated to the primary contact via phone and/or fax.
 Preferred contact: Phone Fax

C. Interconnect Support Services (select 1 option)

Full Interconnect Support Services: Interconnect will provide full reimbursement support including benefit verification, provide prior authorization and appeals support, evaluate patients for eligible financial assistance programs, and enroll patients into adherence services.

Patient Support Services Enrollment Only: My patient is already receiving services from their specialty pharmacy, and I would like them to receive only adherence and education services from Interconnect.

Interim Access Program/Patient Assistance Program: My patient requires evaluation of patient support programs based on their eligibility due to indicating financial concern or coverage through their insurance provider (must select IAP/PAP in Section D).

[ICPT DV04]

D. Prescription and pharmacy information

Prescription for OCALIVA® (obeticholic acid)

5 mg, PO once daily x 30 days, #30 tablets # of refills _____
 10 mg, PO once daily x 30 days, #30 tablets # of refills _____
 Specialty Pharmacy Prescription
 Interim Access Program (IAP)/Patient Assistance Program (PAP) Prescription through AllCare Plus Pharmacy

Select **both** Specialty Pharmacy Prescription and Interim Access Program/PAP checkboxes if you would like your patient to be reviewed for eligibility in the event the patient is experiencing insurance delays or is uninsured (Terms and Conditions apply). Only one pharmacy may receive this prescription to dispense.

Additional considerations: _____

Allergies: _____

Concurrent medications: _____

Pharmacy

Preferred specialty pharmacy: _____

Prescription already sent to pharmacy: Yes No

(Please note that Interconnect's limited specialty pharmacy network includes AcariaHealth™, Accredo Health Group, Inc., AllianceRx Walgreens Pharmacy, CVS Specialty®, CenterWell Specialty Pharmacy, and Optum® Specialty Pharmacy)



Prescriber authorization

I authorize Intercept Pharmaceuticals, Inc. as my designated agent and on behalf of my patient to (1) forward this statement of medical necessity to furnish any information on this form to and recruit necessary patient information from the insurer of above-named patient and (2) forward this prescription, by fax or any means under applicable law, to the pharmacy. I certify that the rationale for prescribing OCALIVA is for a primary diagnosis of ICD-10: K74.3, and I will be supervising the patient's treatment accordingly.

I also certify that I understand OCALIVA is contraindicated in patients with decompensated cirrhosis (e.g., Child-Pugh B or C), a prior decompensation event, with compensated cirrhosis who have evidence of portal hypertension, or with complete biliary obstruction according to the Full Prescribing Information. In addition, I understand that hepatic decompensation and failure, sometimes fatal or resulting in liver transplant, has been reported with OCALIVA treatment in PBC patients with either compensated or decompensated cirrhosis. Understanding this and other information contained in the Full Prescribing Information, I have determined that OCALIVA is appropriate for this patient.

I provide authorization to Interconnect and its affiliates to call, text, or email the patient in regard to securing proper documentation for their services.

Please select **1 option** and sign **only once** below.

	OR	
Prescriber's signature (no stamps; substitution permitted)		Prescriber's signature (no stamps; dispense as written)
_____		_____
Date		Date

Intercept makes no representation that the information will comply with the requirements of any particular payer/insurer. The use of this information does not guarantee payment or that any payment received will cover prescription costs.

Special note: The physician is to comply with their state-specific prescribing requirements, such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance of state-specific requirements could result in outreach to the prescriber.

Please sign and return the completed form and required documentation via fax to [1-855-686-8730]

Please see Important Safety Information on page 4, and visit www.interceptpharma.com/ocaliva-us-prescribing-information for Full Prescribing Information, including Boxed WARNING. Rx only.



Patient name

Date of birth

Patient email

Phone number

I. Patient Authorization to Share Personal Health Information

The Interconnect® Support Services program provides services which vary from patient to patient, such as prescription management, support in securing reimbursement, referrals to patient financial support programs, drug shipment and refills outreach, compliance, and persistency messaging to patients and the patients' physicians, no-cost medication to qualified patients, and other related services in connection with Intercept products and programs (the "Program"). The Program services may change from time to time.

This Authorization will allow the patient's healthcare provider(s) and health insurer(s) to share information with Intercept Pharmaceuticals, Inc. and its affiliates and their employees, including field representatives (collectively, "Intercept"), as well as third-party companies working on Intercept's behalf, for the purposes described in this Authorization.

AUTHORIZATION: By signing this Authorization, I (the patient or the patient's personal representative on behalf of the patient) authorize each of my physicians, pharmacists, and other healthcare providers (collectively, "Healthcare Providers") and each of my health insurers (collectively, "Insurers") to use and/or disclose the personal information described below to Intercept and third parties administering Program services (including any Intercept service providers), for the purposes described in this Authorization. I understand that my pharmacy providers may receive remuneration for disclosing my protected health information and/or for any patient support activities pursuant to this Authorization. My personal health information may be disclosed orally or in writing, including by facsimile, email, and/or through other data transfer means.

My Healthcare Providers and Insurers may disclose to Intercept and other third parties helping to administer Program services (including any Intercept service providers) my personal health information such as: (1) my name, birth date, address, or telephone number; (2) medical records and treatment information; (3) information about my health benefits or health insurance coverage; and (4) financial information about me.

I understand that once my protected health information has been disclosed to Intercept, federal privacy laws may no longer protect the information from further disclosure, but Intercept has agreed to use and disclose my information only for purposes of providing Program services or as indicated in this Authorization.

Intercept may use and/or contact me about my personal health information for the following purposes:

- Contact me by phone, mail, or email to provide information about the Program;
- Verify the accuracy of the information on this form and request additional financial and insurance information;
- Determine my eligibility for the Program and the specific services of the Program, such as financial assistance;
- Facilitate the provision of Program services to me, including delivering Program services such as compliance and persistency messaging and analyzing the impact of the program;
- To send enrollment information to my pharmacy via phone/oral, email, postal mail, or fax;
- For Intercept's medical research and related purposes, including to help develop new products, services, and programs; and/or
- As necessary to comply with applicable laws, including, without limitations, any safety reporting obligations.

I also understand that:

- I do not have to sign this Authorization. My treatment, payment for treatment, insurance enrollment, or eligibility for insurance benefits will not be affected. If I do not sign this Authorization, however, I will not be eligible to receive Program services.
- This Authorization will remain in effect until I am no longer participating in the Program, until the Authorization is required to be terminated pursuant to applicable laws, or until I revoke this Authorization.
- I may revoke (cancel) this Authorization at any time by mailing a letter requesting such cancellation to Interconnect at [P.O. Box 580, Somerville, NJ, 08876] or by emailing Interconnect at [privacyprotection@interceptpharma.com]. If I revoke this Authorization, my Healthcare Providers and Insurers are not permitted to make further disclosures of my personal health information to Intercept, except for disclosures made in reliance on this Authorization. Revocation of this Authorization will not affect Intercept's ability to use or disclose information it has already received. If I revoke this Authorization, I will no longer be able to receive Program services.
- I am entitled to a copy of this signed Authorization.
- Intercept may change or discontinue the Program at any time. Significant changes to the Program will be communicated in a timely manner to all participants of the Program.
- The information disclosed pursuant to this Authorization may be subject to redisclosure by the recipient and may no longer be protected by HIPAA once disclosed.
- I will contact the Program if my financial status or insurance coverage changes.
- Pharmacy may receive financial remuneration from the manufacturer for disclosing PHI to provide services within the authorization

Please sign and return the completed form on Page 1 and required documentation via fax to

[1-855-686-8730]

Please see Important Safety Information on page 6, and visit [www.interceptpharma.com/ocaliva-us-prescribing-information] for Full Prescribing Information and Medication Guide for OCALIVA. Rx only.

Interconnect Savings Program Terms and Conditions

The INTERCONNECT Savings Program ("Program") will provide financial assistance for the out-of-pocket costs for eligible commercially insured patients with a valid prescription, up to a maximum of \$10,000 in assistance per patient per calendar year.

The INTERCONNECT Savings Program is not insurance and is not intended to substitute for insurance.

For the Program, your personal information will be provided to Intercept Pharmaceuticals, Inc. ("Intercept"). This may include your name, address, phone number, email address, and information related to your prescription medication insurance and treatment. This information is needed for Intercept Pharmaceuticals, Inc., and our service providers to enroll you in the Program. We may also use the information you give us to learn more about the patients who use our products and to improve the information we provide to them. Intercept will not share your information with other parties outside of the Program, except where legally required. The information you provide will be processed in accordance with the consent you provided when enrolling in the Program.

The patient must meet the Program requirements every time they use the INTERCONNECT Savings Program. **The Program terms will expire at the end of each calendar year. The Program may change or end for any reason without notice, including within specific states.**

In order to participate in the Program and receive financial assistance, the patient must meet certain eligibility criteria and comply with all the terms and conditions described below:

- **This Program is only available for patients 18 years or older who use commercial or private insurance.**
- To enroll in the Program, the patient (or the patient's legal representative on behalf of the patient, as applicable) must personally complete the enrollment process for the Program. Third-party payers, pharmacy benefit managers, or the agents of either, are prohibited from assisting patients with enrolling in the Program. Any decision to enroll in the Program must be made voluntarily by the patient.
- **This Program is not for patients who use:**
 - **any state or federal government funded health program. Examples of these programs are Medicaid, Medicare, Medigap, Veterans Affairs, Department of Defense, Tricare, the Puerto Rico Government Health Plan, or other federal or state healthcare programs (including any state prescription drug assistance programs); or**
 - **private insurance plans or other health or pharmacy benefit programs that reimburse you for the entire cost of your prescription drugs.**
 - Uninsured and cash-paying patients are not eligible to enroll in the Program.
- Patients who begin receiving prescription benefits from any state or federal government funded health program at any time must notify Intercept of this fact by contacting [CONTACT] and will no longer be eligible for this Program.
- The Program is limited to one per person and is not transferable. No substitutions are permitted. This Program is offered to, and intended for the sole benefit of, eligible patients and may not be utilized for the benefit of third parties, including, without limitation, third-party payers, pharmacy benefit managers, or the agents of either.
- The savings received under this Program must be deducted from any reimbursement request submitted to the patient's insurance plan, either directly or on behalf of the patient.
- The patient and the pharmacist each must report the patient's receipt of financial assistance under this program as required by any insurer, health plan, or other third-party payer.
- Void where prohibited by law, taxed, or restricted.
- This Program offer may not be used with any other coupon, discount, prescription savings card, free trial, or other offer (including, without limitation, any program offered by a third-party payer or pharmacy benefit manager, or an agent of either, that adjusts patient cost-sharing obligations).

This Program offer is valid only to eligible residents of the United States and Puerto Rico and void where prohibited, taxed, or limited by law.

You may end your participation in the INTERCONNECT Savings Program at any time by calling [844-622-4278.]

You may end your participation in the INTERCONNECT Savings Program at any time by calling [1-844-622-4278].

Please see Important Safety Information on page 6, and visit [www.interceptpharma.com/ocaliva-us-prescribing-information] for Full Prescribing Information and Medication Guide for OCALIVA. Rx only.

HCP INDICATION AND IMPORTANT SAFETY INFORMATION

INDICATION

OCALIVA® (obeticholic acid), a farnesoid X receptor (FXR) agonist, is indicated for the treatment of adult patients with primary biliary cholangitis (PBC)

- without cirrhosis or
- with compensated cirrhosis who do not have evidence of portal hypertension,

either in combination with ursodeoxycholic acid (UDCA) with an inadequate response to UDCA or as monotherapy in patients unable to tolerate UDCA.

This indication is approved under accelerated approval based on a reduction in alkaline phosphatase (ALP). An improvement in survival or disease-related symptoms has not been established. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

IMPORTANT SAFETY INFORMATION

WARNING: HEPATIC DECOMPENSATION AND FAILURE IN PRIMARY BILIARY CHOLANGITIS PATIENTS WITH CIRRHOSIS

- **Hepatic decompensation and failure, sometimes fatal or resulting in liver transplant, have been reported with OCALIVA treatment in primary biliary cholangitis (PBC) patients with either compensated or decompensated cirrhosis.**
- **OCALIVA is contraindicated in PBC patients with decompensated cirrhosis, a prior decompensation event, or with compensated cirrhosis who have evidence of portal hypertension.**
- **Permanently discontinue OCALIVA in patients who develop laboratory or clinical evidence of hepatic decompensation; have compensated cirrhosis and develop evidence of portal hypertension; or experience clinically significant hepatic adverse reactions while on treatment.**

Contraindications

OCALIVA is contraindicated in patients with:

- decompensated cirrhosis (e.g., Child-Pugh Class B or C) or a prior decompensation event.
- compensated cirrhosis who have evidence of portal hypertension (e.g., ascites, gastroesophageal varices, persistent thrombocytopenia).
- complete biliary obstruction.

Warnings and Precautions

Hepatic Decompensation and Failure in PBC Patients with Cirrhosis

Hepatic decompensation and failure, sometimes fatal or resulting in liver transplant, have been reported with OCALIVA treatment in PBC patients with cirrhosis, either compensated or decompensated. Among postmarketing cases reporting it, median time to hepatic decompensation (e.g., new onset ascites) was 4 months for patients with compensated cirrhosis; median time to a new decompensation event (e.g., hepatic encephalopathy) was 2.5 months for patients with decompensated cirrhosis. Some of these cases occurred in patients with decompensated cirrhosis when they were treated with higher than the recommended dosage for that patient population; however, cases of hepatic decompensation and failure have continued to be reported in patients with decompensated cirrhosis even when they received the recommended dosage.

Hepatotoxicity was observed in the OCALIVA clinical trials. A dose-response relationship was observed for the occurrence of hepatic adverse reactions including jaundice, worsening ascites, and primary biliary cholangitis flare with dosages of OCALIVA of 10 mg once daily to 50 mg once daily (up to 5-times the highest recommended dosage), as early as one month after starting treatment with OCALIVA in two 3-month, placebo-controlled clinical trials in patients with primarily early stage PBC.

Routinely monitor patients for progression of PBC, including hepatic adverse reactions, with laboratory and clinical assessments to determine whether drug discontinuation is needed. Closely monitor patients with compensated cirrhosis, concomitant hepatic disease (e.g., autoimmune hepatitis, alcoholic liver disease), and/or with severe intercurrent illness for new evidence of portal hypertension (e.g., ascites, gastroesophageal varices, persistent thrombocytopenia) or increases above the upper limit of normal in total bilirubin, direct bilirubin, or prothrombin time to determine whether drug discontinuation is needed. Permanently discontinue OCALIVA in patients who develop laboratory or clinical evidence of hepatic decompensation (e.g., ascites, jaundice, variceal bleeding, hepatic encephalopathy), have compensated cirrhosis and develop evidence of portal hypertension (e.g., ascites, gastroesophageal varices, persistent thrombocytopenia), experience clinically significant hepatic adverse reactions, or develop complete biliary obstruction. If severe intercurrent illness occurs, interrupt treatment with OCALIVA and monitor the patient's liver function. After resolution of the intercurrent illness, consider the potential risks and benefits of restarting OCALIVA treatment.

Severe Pruritus

Severe pruritus was reported in 23% of patients in the OCALIVA 10 mg arm, 19% of patients in the OCALIVA titration arm, and 7% of patients in the placebo arm in a 12-month double-blind randomized controlled clinical trial of 216 patients. Severe pruritus was defined as intense or widespread itching, interfering with activities of daily living, or causing severe sleep disturbance, or intolerable discomfort, and typically requiring medical interventions. Consider

clinical evaluation of patients with new onset or worsening severe pruritus. Management strategies include the addition of bile acid binding resins or antihistamines, OCALIVA dosage reduction, and/or temporary interruption of OCALIVA dosing.

Reduction in HDL-C

Patients with PBC generally exhibit hyperlipidemia characterized by a significant elevation in total cholesterol primarily due to increased levels of high-density lipoprotein-cholesterol (HDL-C). Dose-dependent reductions from baseline in mean HDL-C levels were observed at 2 weeks in OCALIVA-treated patients, 20% and 9% in the 10 mg and titration arms, respectively, compared to 2% in the placebo arm. Monitor patients for changes in serum lipid levels during treatment. For patients who do not respond to OCALIVA after 1 year at the highest recommended dosage that can be tolerated (maximum of 10 mg once daily), and who experience a reduction in HDL-C, weigh the potential risks against the benefits of continuing treatment.

Adverse Reactions

The most common adverse reactions ($\geq 5\%$) are: pruritus, fatigue, abdominal pain and discomfort, rash, oropharyngeal pain, dizziness, constipation, arthralgia, thyroid function abnormality, and eczema.

Drug Interactions

• Bile Acid Binding Resins

Bile acid binding resins such as cholestyramine, colestipol, or colesevelam adsorb and reduce bile acid absorption and may reduce the absorption, systemic exposure, and efficacy of OCALIVA. If taking a bile acid binding resin, take OCALIVA at least 4 hours before or 4 hours after taking the bile acid binding resin, or at as great an interval as possible.

• Warfarin

The International Normalized Ratio (INR) decreased following coadministration of warfarin and OCALIVA. Monitor INR and adjust the dose of warfarin, as needed, to maintain the target INR range when co-administering OCALIVA and warfarin.

• CYP1A2 Substrates with Narrow Therapeutic Index

Obeticholic acid may increase the exposure to concomitant drugs that are CYP1A2 substrates. Therapeutic monitoring of CYP1A2 substrates with a narrow therapeutic index (e.g., theophylline and tizanidine) is recommended when co-administered with OCALIVA.

• Inhibitors of Bile Salt Efflux Pump

Avoid concomitant use of inhibitors of the bile salt efflux pump (BSEP) such as cyclosporine. Concomitant medications that inhibit canalicular membrane bile acid transporters such as the BSEP may exacerbate accumulation of conjugated bile salts including taurine conjugate of obeticholic acid in the liver and result in clinical symptoms. If concomitant use is deemed necessary, monitor serum transaminases and bilirubin.

Visit www.interceptpharma.com/ocaliva-us-prescribing-information for Full Prescribing Information, including Boxed WARNING.

To report SUSPECTED ADVERSE REACTIONS, contact Intercept Pharmaceuticals, Inc. at 1-844-782-ICPT or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

You are eligible for this program if:

- You do not have any prescription drug coverage for OCALIVA® (obeticholic acid)
- You are an adult (18 years and older) diagnosed with primary biliary cholangitis
- You are a US citizen or permanent resident of the United States or Puerto Rico
- Your annual gross household income is at or below 500% of the Federal Poverty Level for all family sizes. Please visit the US Department of Health & Human Services, Office of the Assistant Secretary for Planning and Evaluation, to view the current Federal Poverty Guidelines at <https://aspe.hhs.gov/poverty-guidelines>

Patient information (to be completed by the patient)

_____ / / _____ Male Female
 First name Last name Date of birth

 Address City

_____ Preferred contact: _____ Email _____ Home phone _____ Cell phone _____
 State ZIP

 Email address Home phone Cell phone

Are you a US citizen? YES NO If no, are you a permanent resident of the United States? YES NO

Total household income \$ _____ No. of people in your household: _____

Do you have private prescription insurance coverage? YES NO

Have you enrolled in Medicaid? YES NO

Are you enrolled in Medicare Part A and/or Part B? YES NO Medicare ID no. (if applicable): _____

Ship OCALIVA to: Patient's home HCP's office

Patient declaration

AUTHORIZATION: I know that to qualify for free medicine my household gross income must be at or below 500% of the Federal Poverty Level, and I certify that the patient financial information I have provided is correct. I certify I have no health plan coverage for OCALIVA; this includes Medicare, Medicaid, or other public programs. I do not have the resources to pay for OCALIVA. I agree to provide Interconnect proof of my income, if requested. I agree that if my certification about my income is false, I will reimburse Intercept Pharmaceuticals, Inc.

➔ _____
 Patient's signature Date

Statement of Medical Necessity

To the best of my knowledge, this patient has no coverage (including Medicare, Medicaid, or other Federal healthcare programs) for OCALIVA. I certify that, in my medical judgment, OCALIVA is medically necessary for this patient, and that I will be supervising this patient's treatment.

➔ _____
 Prescriber's signature Date

Please sign and fax the completed form and required documentation to:
[1-855-686-8730]
 Additional documentation attached

Please see Important Safety Information on page 6, and visit www.interceptpharma.com/ocaliva-us-prescribing-information for Full Prescribing Information and Medication Guide for OCALIVA. Rx only.

What is OCALIVA® (obeticholic acid)?

OCALIVA is a prescription medicine used to treat primary biliary cholangitis (PBC) in combination with ursodeoxycholic acid (UDCA) in adults who have not responded well enough to UDCA, or alone for adults who cannot tolerate UDCA. It is not known if taking OCALIVA will improve your chance of survival or improve your symptoms of PBC. It is not known if OCALIVA is safe and effective in children.

What is the most important information I should know about OCALIVA?

OCALIVA may cause serious side effects including:

Worsening of liver problems or liver failure, in some cases leading to liver transplant or death, has happened in people with primary biliary cholangitis (PBC) with liver cirrhosis when taking OCALIVA.

Before you start OCALIVA, and during your treatment with OCALIVA, your healthcare provider will do tests to check your liver. These tests will help your healthcare provider decide if it is safe for you to start taking OCALIVA and safe for you to continue taking OCALIVA.

Tell your healthcare provider right away if you have any of the following symptoms of worsening liver problems during treatment with OCALIVA:

- Swelling of your stomach-area from a build-up of fluid; yellowing of your skin or the whites of your eyes; black, tarry, or bloody stools; coughing up or vomiting blood, or your vomit looks like "coffee grounds"; mental changes such as confusion, sleepier than usual or harder to wake up, slurred speech, mood swings, or changes in personality.

Tell your healthcare provider right away if you have any of the following symptoms during treatment with OCALIVA and they are severe or do not go away:

- Stomach-area pain; nausea, vomiting, or diarrhea; loss of appetite or weight loss; new or worsening fatigue; weakness; fever and chills; light-headedness; less frequent urination

Who should not take OCALIVA?

Do not take OCALIVA if you:

- have PBC with liver cirrhosis with symptoms such as fluid in the stomach-area or confusion (decompensated liver cirrhosis) or with abnormalities in certain tests that check your liver.
- have a complete blockage of the bile ducts in your liver or gallbladder.

What are the possible side effects of OCALIVA?

OCALIVA may cause serious side effects, including:

- See "What is the most important information I should know about OCALIVA?"
- **Severe Itching (pruritus).** Itching is a common side effect and can sometimes become severe (intense itching or itching over much of your body). Severe itching can cause discomfort, problems sleeping, and problems doing daily activities and usually needs to be treated. Tell your healthcare provider if you get severe itching or if your itching gets worse.
- **Lower HDL-C** ("good" cholesterol). OCALIVA can lower high levels of HDL-C. Your healthcare provider will check your cholesterol levels during treatment with OCALIVA.

The most common side effects of OCALIVA include: tiredness; stomach pain and discomfort; rash; joint pain; mouth and throat pain; dizziness; constipation; swelling in your hands, ankles, or feet; fast or irregular heartbeat; fever; changes in how your thyroid gland works; dryness, irritation, redness, crusting or drainage of the skin (eczema).

These are not all the possible side effects of OCALIVA. Call your doctor for medical advice about side effects.

What should I tell my healthcare provider before taking OCALIVA?

Before taking OCALIVA, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if OCALIVA will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OCALIVA passes into your breastmilk. Talk with your healthcare provider about the best way to feed your baby if you take OCALIVA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. OCALIVA can affect the way certain medicines work. Certain other medicines may affect the way OCALIVA works.

The risk information provided here is not complete. To learn more, please talk to your healthcare provider.

Visit [www.interceptpharma.com/ocaliva-us-prescribing-information] for Full Prescribing Information and Medication Guide for OCALIVA.

Available by prescription only.

To report negative side effects of OCALIVA, please contact Intercept Pharmaceuticals, Inc. at 1-844-782-ICPT or you may report to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.