Date Shipment Needed:	Ship To: □Patient □Prescriber
□ Nursing needed; □Training needed ► All the supplies incl	uding syringes and needles will be dispensed if needed.

INJECTAFER REFERRAL FORM

PATIENT INFORMATION											
Patient Name:			DOB:	Ç	Sex: □M □F	Weight:		□lbs. □kg.			
SSN	N:	Phone:	Allergies:								
Add	ress:		-	City:		State:	Zip:				
Emergency Contact: Phone:			Phone:			☐ Please attach demographic information					
PRE	ESCRIBER INFORMATION										
Prescriber:			NPI:	DEA:		State Lic:					
	pervising Physician:			Practice N	Name:	T					
	dress:	T-		City:	2 1 1	State:	Zip:				
	one:	Fax:		Key Office	e Contact:		Phone:				
DIAGNOSIS INFORMATION / MEDICAL ASSESMENT Primary Diagnosis DDF0 0 large deficiency approximation to blood loss. DDF0 0 Other land deficiency. DDF0 1 Approximation in changing in changi											
Primary Diagnosis: □D50.0 Iron deficiency anemia due to blood loss □D50.8 Other Iron deficiency □D63.1Anemia in chronic kidney disease □Other:											
	Does the patient have non-dialysis dependent chronic kidney disease? □Yes □No										
•	Is the patient currently on dialysis? ☐Yes ☐No										
•	■ Does the patient experience an intolerance to an an oral iron or an unsatisfactory response to oral iron? □Yes □No Medication(s):										
	Is patient currently on therapy? Yes No Medication(s):										
•	Authinistering as undiluted slow intraversous Fusin: 4 Fes 4No										
■ Administering as an Infusion? □Yes □No Infusion supplies:											
INSURANCE INFORMATION											
□ Please attach front and back of patient's insurance card (medical and prescription)											
	PAY CARD ENROLLMENT	construction Construction									
□ Please check if enrolling in copay card Copay ID: PRESCRIPTION INFORMATION											
PKE	SCRIPTION INFORMATION										
□lnj	jectafer® ferric carboxymaltos										
□<50 kg 15 mg/kg, give Injectafer in 2 doses separated by at least 7 days and each						ials	QTY:	Refills:			
□>50 kg, give Injectafer in 2 doses of 750 mg doses separated by at least 7 day			it least / days, /	750 mg/mL vials				QTY:	Refills:		
		ig of from per course									
□Infusion Supplies Required □Sterile 0.9% Sodium Chlorida, USP, use saline to dilute Injectafer to 2 to 4 mg of iron per mL for infusion (dilute up to 750 mg of iron in no more than 250 mL of strelie 0.9% sodium chloride injection or infusion bag) □Other supplies:						0=1/ =00 /	5 5"				
								QTY: <u>500 mL</u>	Refills:		
			injection or iniusion pay)				QTY:	Refills:			
	- outroi supplies.								i tollilo.		